

Ultraslow-Spreading Ridges: Rapid Paradigm Changes

Second Final submission to Oceanography v1.2 - 12.29.2006

Jonathan E. Snow¹ and Henrietta N. Edmonds²

Jonathan E. Snow (jesnow@uh.edu) is Assistant Professor, University of Houston, Department of Geosciences, Houston, TX, USA. **Henrietta N. Edmonds** is Assistant Professor, The University of Texas at Austin, Marine Science Institute, Port Aransas, TX, USA.

Ultraslow spreading ridges (<20mm/yr full rate) represent a major departure from the style of crustal construction seen in the rest of the ocean basins. Since the 1960's, fast and slow spreading ridges of the Pacific and Atlantic, combined with the ophiolite model of crustal structure, have defined the conceptual context for the structural, tectonic and hydrothermal architecture of mid-ocean ridges. Over the last 15 years, studies of ultraslow spreading ridges have presented researchers with anomalies not explained by the standard model of crustal formation, and in recent years they have become recognized as a class unto themselves. Their "anomalous" characteristics in fact provide key information about many of the underlying processes that govern crustal accretion at all spreading rates. The ultraslow ridges include the SW Indian Ridge (between Africa and Antarctica) the Gakkel Ridge (bisects the Arctic Ocean) and several smaller ridges (Figure 1).




Figure 1. Global plate boundaries and oceanic spreading ridge segments (Bird, 2003). Green indicates fast spreading ridges (full spreading rate ≥ 60 mm/yr). Red indicates ultraslow spreading ridges. All other ridge segments are indicated in yellow. GR = Gakkel Ridge; LT = Lena Trough; KR = Knipovich Ridge; MR = Mohns Ridge; CT = Cayman Trough; AAR = America-Antarctic Ridge; SWIR = Southwest Indian Ridge.

Since the advent of plate tectonic theory, mid-ocean ridges have been classified based on their structural, morphological and volcanologic characteristics into two major types based largely on their spreading rate: “fast” (>60 mm yr⁻¹) and “slow” (<60 mm yr⁻¹ full rate); an “intermediate” type is often placed between them. Both types of ridge share certain characteristics: (1) they have roughly the same crustal thickness (6-7 km; Figure 2), (2) in plan view they have a characteristic stair step geometry of volcanic rifts separated by perpendicular transform offsets, and (3) they generate a characteristic outcrop pattern of elongate fault-bounded abyssal hills trending normal to the spreading direction. Their differences lie primarily in their across-axis morphology: Fast spreading ridges have an axial rise with a very narrow summit graben that is the locus for most volcanic

and tectonic activity, whereas slow spreading ridges have rugged rift mountains enclosing a broad axial valley. Fast spreading ridges tend to be dominated by volcanism, slow spreading ridges more by tectonics. This distinction was widely accepted for many years as a basic principle by the worldwide geologic community. It was taught as a fundamental characteristic of the geology of the sea floor to undergraduates, and its causes were debated as one of the not-completely-understood corners of plate tectonic theory.




Figure 2: Crustal thickness as a function of spreading rate. Crustal thickness was determined by seismic refraction, modified after (Bown and White, 1994). Normal oceanic crust has a mean thickness of 6 km at all spreading rates above 20 mm yr⁻¹.

The characteristics of both slow- and fast-spreading ridges fit well with the ophiolite model for the formation of the ocean crust, which entered the geological canon at the

1972 Penrose Conference on Ophiolites and Ocean Crust (Penrose Participants, 1973). This model calls for a layered structure of pillow basalts, sheeted dikes, gabbro and mantle (see Figure 3), in a thickness and proportion consistent with the seismic structure of both fast and slow spreading crust. These characteristics, verified by geological mapping on land in ophiolites, have been an important cornerstone of plate tectonic theory for the past 35 years, and have continuously proven useful in helping us understand the most inaccessible parts of the Earth's crust. (Nicolas, 1995).

Problems with the Penrose Model at the Slowest Spreading Rates

Outcrop of mantle ultramafic rocks on the ocean floor was first described at the slow spreading Mid-Atlantic Ridge (Aumento and Loubat, 1970). In the Penrose model, a 6 km layer of basaltic rock covers the mantle, thus ultramafic rocks should be rare. Their emplacement to the ocean floor requires mechanisms that would seem implausible, such as faults with a minimum of 6 km displacement (though such faults were in fact later found), or serpentinite diapirism when the serpentinites themselves have densities hardly less than the basaltic rocks they must rise through. Also, there is no indication how water might penetrate through kilometers of ocean crust to create serpentinites in the mantle. Early on, these problems were explained by the idea that great transform faults (Morgan, 1968) offsetting ridge segments provided a pathway for water to enter the mantle, and for serpentine diapirs to rise to the surface. Most abyssal peridotites were recovered from the walls of large-offset transform faults, or near them. On the Southwest Indian Ridge (SWIR), peridotites were even more common (Dick, 1989; Engel and Fisher, 1975).




Figure 3: Model of oceanic crustal structure (Nicolas, 1995). The Harzburgite type describes crust similar to “Normal” Penrose-style mid-ocean ridge crust. The origin of the Lherzolite type was debated for many years, but is now correlated with nonvolcanic rifted margins and to ultraslow spreading ridges.

By the mid-1990’s, unusual ridge segments were also found on the SW Indian Ridge that were anomalously deep and not perpendicular to the spreading direction, as Penrose-style volcanic rifts are supposed to be. Neither were they transform faults, which are geometrically required to be parallel to the spreading direction – the walls of these rifts instead trend at a highly oblique angle to the prevailing spreading direction. These unusual oblique rift segments frequently contained mantle peridotites, as well as basalts of unusual composition. Although these types of basalt were already known from ocean islands and ridges near major hotspots, their presence in these anomalous rifts, often called “leaky transforms,” could not easily be reconciled with prevailing ideas about

crustal accretion or the generation of oceanic basaltic magmas (Cannat et al., 1999; le Roex et al., 1992; Patriat et al., 1997)

A third major change to ideas of crustal accretion came through discoveries at ridge-transform intersections. Dredging results showed that the elevated inside corner high sections of transform faults contained abundant rocks from the lower crust and upper mantle. While the faults bounding the inside corner high were large, with as much as 500m of obvious vertical displacement, this was nowhere near enough to bring up mantle rocks from beneath a full section of oceanic crust (Dick, 1989; Karson, 1983). Subsequently, improved bathymetric imaging showed obvious signs that this faulting was along many 10's of kilometers of displacement (Cann et al., 1997; Tucholke et al., 1998). Nearby dredging of anomalously *deep* regions of the ridge (so-called zero-offset fracture zones, or non-transform discontinuities) also recovered lower crustal and upper mantle rocks, far from the effects of transforms (Cannat et al., 1995). At this point, one was forced to wonder, even on slow spreading ridges, just how much of the crust could be considered “normal”.

Almost from the beginning, the observation of a constant crustal thickness across the entire range of spreading rates (cf Figure 2) began to unravel as well. While it remained true at most spreading ridges that the seismically determined crustal thickness was nearly constant, at the slowest spreading rates, notably in a seismic study done through the ice of the Arctic ocean, the ocean crust seemed to be dramatically thinner than along the rest of the global mid-ocean ridge system (Figure 2). So thin in fact, that the very concept of a “crust” had to be called into question, as the seismic structures

found could easily be satisfied by a thin layer of serpentinite overlying bare mantle (Bown and White, 1994; Jackson et al., 1982; Reid and Jackson, 1981).

Evidence from the frozen North: AMORE 2001

A consensus thus evolved during the 1990's that while the Penrose model still held for most mid-ocean ridge systems, the very slowest spreading ridges were quite different, and might thus illustrate extreme forms of the fundamental geodynamic forcing functions driving mid-ocean ridge construction everywhere. A major effort then evolved to investigate the world's slowest spreading mid-ocean ridge – the Gakkel Ridge in the Arctic Ocean (InterRidge-Arctic-Working-Group, 1998). Gakkel Ridge is the northern continuation of the Mid-Atlantic Ridge, and extends over 2000 km from the NE tip of Greenland to the Laptev Sea continental shelf in Siberia. The full spreading rate varies from about 14mm yr^{-1} at the Greenland end to less than 8 mm yr^{-1} at its Laptev Sea termination (Reid and Jackson, 1981; Vogt et al., 1979). Thus, at its fastest end, the Gakkel Ridge is slower than any other mid-ocean ridge. Pack ice cover had thus far prevented Arctic hard rock dredge operations, until in 1999 the German PFS Polarstern, recovered the first peridotite, basalt and hydrothermal rocks from the Lena Trough, just to the south (Snow et al., 2001b). At the same time, a US Navy-civilian science cooperation used civilian instruments on a US nuclear submarine to produce the first bathymetric and sidescan sonar images of Gakkel Ridge (Coakley and Cochran, 1998; Cochran, 2003; Edwards et al., 2001).




Figure 4: Gakkel Ridge Bathymetry (Jokat et al., 2003, Fig. 1, modified) based on AMORE 2001 and IBCAO data sets. Full spreading rates in each identified section are given in mm yr⁻¹. Red lines show magnetic lineation picks.

The international mapping and sampling program that left Tromsø, Norway in July of 2001, AMORE 2001, was an unprecedented success (Edmonds et al., 2003; Jokat et al., 2003; Michael et al., 2003). Encountering good ice conditions, scientists aboard PFS *Polarstern* (Alfred Wegener Institute, Bremerhaven, Germany) and the new US Coast Guard Cutter *Healy* were able to create a detailed bathymetric map of much of the rift valley of Gakkel Ridge (Figure 4), carry out over 200 rock sampling stations, a dozen seismic crustal thickness measurements, and continuous seismic reflection seismic transects across both the Nansen and Amundsen basins. The two ships reached the North

Pole together on September 6, 2001 (Jokat et al., 2003; Michael et al., 2003). In 2004 Polarstern mapped and sampled the Lena Trough and returned to Gakkel Ridge for additional mapping and sampling (Snow and Petrology-Group-ARK-XX-2, 2006).

A “new” class of mid-ocean ridge

The new observations made along Gakkel Ridge were confusing. There was significantly more volcanism than had been expected. The western part of Gakkel Ridge was magmatically robust, despite its relatively thin crust, with no outcrops of mantle peridotite. On either side of this western volcanic zone, however, were long rift segments where apparently no magmatic crustal construction occurred at all (Michael et al., 2003; Snow and Petrology Group, ARK XX/2, 2006). To the east, however, volcanism *increased* in discrete volcanic centers separated by amagmatic basins.

The SW Indian and Gakkel Ridge together thus required a new entry in the table of mid-ocean ridge types (Dick et al., 2003): At ultraslow spreading ridges, the seismic crust is thinner (~1-4 km) than at other ridges (see Figure 2) (Bown and White, 1994; Jokat et al., 2003; Reid and Jackson, 1981). The changeover from slow to ultraslow spreading characteristics is not a strict function of spreading rate. Rather, a number of factors can influence the style of crustal accretion: Plate boundary geometry, mantle magmatic productivity and mantle potential temperature all seem to play a role (Dick et al., 2003).

Ultraslow spreading ridges uniquely possess amagmatic rifts that expose mantle peridotite directly on the sea floor, with only scattered basalt and gabbro (Dick et al., 2003). Along with magmatic rifts, transforms and subduction zones, amagmatic rifts form a new fourth class of plate boundary structure. Figure 5 shows an amagmatic segment on

the SW Indian Ridge (Dick et al., 2003); they also occur on the Gakkel Ridge (Michael et al., 2003) and Lena Trough (Snow and Petrology Group, ARK XX/2, 2006). Unlike magmatic segments, they have mantle peridotite walls formed by long sloping scarps or irregular uplifts rather than the basaltic block normal faulting of Penrose crust (Dick et al., 2003; Michael et al., 2003; Okino et al., 2002). Amagmatic segments can also assume any orientation to the spreading direction, sometimes forming oblique rifts (Dick et al., 2003; Snow et al., 2001b), and can produce a unique “smooth” sea floor (Cannat et al., 2006). Magmatic accretion at ultraslow spreading rates can either be robustly magmatic, as at the orthogonal supersegment of the SW Indian Ridge (Dick et al., 2003) or the western Gakkel Ridge (Michael et al., 2003) or it may be confined to highly focused short magmatic segments (producing crust with orthogonal structures) linked by amagmatic segments. Where the plate boundary runs oblique to the direction of spreading, the amagmatic style of spreading is enhanced, such that oblique amagmatic segments can form at higher spreading rates than otherwise possible (Dick et al., 2003).




Figure 5: Oblique amagmatic accretion on the SW Indian Ridge (Dick et al., 2003). Filled circles show sampling points, giving rock type coded by color as shown in key. Note that the spreading direction (red arrows) is not at a right angle to the strike of the rift. Filled circles show sampling points, giving rock types as shown in the key. This type of spreading, far from being anomalous, is characteristic of ultraslow spreading ridges.

The more variable and alkalic nature of basalts from ultraslow spreading ridges (le Roex et al., 1992; Meyzen et al., 2003; Snow et al., 2001b) can be explained by an overall lower degree of mantle partial melting in the mantle, and the presence of mantle geochemical components not observed at slow and fast spreading ridges (Hart, 1984). Mantle peridotites from ultraslow spreading ridges afford a unique, nearly direct access to the Earth's mantle, and have made possible the conclusion that there is primary mineralogic variability in the upper mantle that affects the process of partial melting (Dick et al., 1984), that mid-ocean ridge melting occurs by a process of near-fractional melting (Johnson et al., 1990), and that deep melts formed in the presence of garnet are a component of mid-ocean ridge basalt (Hellebrand and Snow, 2003).

Insights on continental rifting and ophiolites

Ultraslow spreading ridges have increasingly become a model for understanding some of the more puzzling aspects of ophiolites and continental rifts. The Penrose model had always met with difficulty in explaining some aspects of the tectonic setting and structure of ophiolites. Far too many ophiolites (the “Lherzolite type” in Figure 3 (Nicolas, 1995)) did not conform to the 6km thick crust observed nearly everywhere in the oceans. The proponents of the ophiolite paradigm concluded that those incomplete ophiolites had become “dismembered” during the process of obduction onto the continent. Among the incomplete Lherzolite type ophiolites are the original ones where the association of volcanic rocks, serpentinite and deep sea sediment was first recognized and called ‘ophiolite’ in 1905 (Steinmann, 1905). These contained little middle crust, consisting largely of a thin layer of pillow basalt erupted directly on peridotite and covered by deep-sea sediments. Harry Hess, a Princeton mineralogist and early mentor to the founders of plate tectonic theory first recognized their significance as the trace of ancient plate boundaries (Hess, 1962). His idea, however, that the ocean floor consisted entirely of serpentinite was ultimately displaced by the Penrose ophiolite model shown in Figure 3.

In the 1990’s however, the Ocean Drilling Program completed a series of deep sea drill holes off the coast of Portugal that showed the existence of serpentinite-floored oceanic crust on the European continental margin (Whitmarsh et al., 1996; Whitmarsh et al., 2001; Whitmarsh et al., 1993). At the same time, the sedimentary sequences the Western Alps were recognized also as the margin of a rifted continent (Froitzheim and Manatschal, 1996; Manatschal, 2004). The basement of this fossil continental rifted margin changes, just as on the present day Iberia Margin, from continental basement to mantle

rocks, sometimes with a thin cover of basalt, and covered by deep sea sediments. The rock types, the structures and ultimately the mechanisms of exhumation of deep crustal and mantle rocks in both of these locations bear striking similarity to that occurring today at ultraslow spreading ridges. Thus, on some continental margins, the rifting that creates new ocean basins occurs in an amagmatic “Hess-type” manner analogous to ultraslow mid-ocean ridges. This combination of Alpine rock types was exactly the one that Harry Hess had noticed the tectonic significance of 50 years ago, and in fact the Western Alpine outcrops involved were the same ones where Steinmann originally described his unique rock association 100 years ago (Steinmann, 1905).

Hydrothermal activity on ultraslow spreading ridges

Ultraslow spreading ridges have also provided surprises and new insights in hydrothermal vent research. Before the first high-temperature venting was observed on the Mid-Atlantic Ridge (Rona et al., 1986) many scientists had been skeptical whether even slow-spreading ridges were capable of supporting such venting due to their colder thermal structure. By the 1990’s a linear relationship was proposed between hydrothermal venting and spreading rate, largely based on results from the faster-spreading ridges (Baker, 1996; Baker et al., 1994). German and Parsons (1998), in extending this work to the slow-spreading Mid-Atlantic Ridge, pointed out that the length scale over which survey results are considered may lead to substantial variability, particularly on slower-spreading ridges, where tectonic fabric plays a stronger role in controlling hydrothermal circulation, and can limit the dispersal of hydrothermal plumes. All of these studies predicted that hydrothermal activity would be very low at ultraslow spreading rates, and the perception persisted that the occurrence of high temperature venting might drop to near

nothing. For the Gakkel Ridge, for example, site vent frequency was predicted to be on the order of one site per 200-300 km.




Figure 6. Summary of known hydrothermal activity on ultraslow spreading ridges. a. Map of the Arctic ridges indicating the area thoroughly surveyed on the AMORE 2001 cruise (box), vent sites identified on AMORE (red stars), and vent sites identified on other cruises (purple stars: Lena Trough, Snow et al. (2001); Knipovich Ridge, Connelly and German (2002); Mohs Ridge., Pedersen et al. (2005)). b. Map of the Southwest Indian Ridge indicating the three areas that have been systematically surveyed for hydrothermal plumes (boxes) and the locations of identified plumes (red stars). For the latter we have indicated all possible plume sites; estimates range from 3 to 8 plumes in the 10-23°E survey area.

It is thus ironic that ultraslow spreading ridges have now been surveyed more thoroughly for hydrothermal activity than most other spreading rate classes (Baker and German, 2004). This is the result of the coincidence between the beginning of the international ridge community's focus on ultraslow spreading ridges and the development of strategies for systematic large scale mapping of hydrothermal plumes e.g. (Baker and Milburn, 1997). Such mapping relies on the plumes' optical backscatter, a measure of suspended particles in the water column that result from the mixture of metal-rich high temperature hydrothermal fluid and seawater. The deployment of optical sensors during rock sampling and deep towed geophysical operations enables much more efficient detection of plumes than the earlier "spot" sampling by CTD. Three sections of ultraslow spreading ridges, totaling ~2250 km, have now been thoroughly surveyed (Figure 6), including the SWIR from 58° to 60°E and 63° to 66°E (German et al., 1998), the SWIR from 10° to 23°E (Bach et al., 2002; Baker et al., 2004), and the Gakkel Ridge from 8°W to 85°E (Edmonds et al., 2003).

The results of these surveys, along with isolated discoveries on the Mohns and Knipovich Ridges (Connelly and German, 2002; Pedersen et al., 2005) and in the Lena Trough (Snow et al., 2001a) (Figure 5), demonstrate that high-temperature hydrothermal venting is ubiquitous on ultraslow-spreading ridges. The three densely surveyed sections yield a site frequency of about one site per 100 km. Recently, normalization of site frequencies to the ridge magma production rate has suggested that ultraslow-spreading ridges may be two to four times as efficient as fast spreading ridges at converting the available magmatic heat into vent fields (Baker et al., 2004; Baker and German, 2004). With the exception of the Mohns Ridge sites (Pedersen et al., 2005), which are located in

about 500 m water depth, all of the sites are in deep water and none have yet been surveyed or sampled with deep submergence technology.

Ultraslow-spreading ridges may have even more hydrothermal activity than outlined above. Optical plume mapping does not detect some forms of hydrothermal activity likely to be more common on slow and ultraslow spreading ridges. Hydrothermal circulation in peridotite-hosted systems is driven by the exothermic serpentinization reactions, rather than by magmatic heat. This results in colder and less metal-rich vent fluids than “traditional” basalt-hosted systems. A case in point is the 40-75°C fluid temperatures at the Lost City ultramafic hydrothermal field in the Atlantic (Kelley et al., 2001). The only water column signal that ultramafic-hosted systems might consistently generate is elevated methane, but as yet there is no reliable, robust sensor for *in-situ* detection of methane at the expected levels. Such “non-traditional” hydrothermal systems may be especially important because of their unique contributions to global geochemical cycles. High-temperature systems that involve magmatic heat as well as ultramafic rocks (such as Rainbow and Logatchev on the northern MAR) also exhibit elevated precious metal concentrations and the abiogenic production of complex (multi-carbon) organic molecules. Furthermore, hydrothermal systems on slow- and ultraslow- spreading ridges may be active for longer periods of time and lead to the deposition of larger ore bodies due to their localization on long-lived fault structures.

Ultraslow spreading ridges occupy regions of particular interest for vent biogeographic studies (Tyler and Young, 2003; Van Dover et al., 2002). The deep Arctic ridges are isolated from the rest of the mid-ocean ridge system due to the presence of Iceland astride the Mid-Atlantic Ridge, and to the shallow connections between the Arctic

Ocean and the Pacific (through the Bering Strait) and between the Arctic and the Atlantic (through the passages of the Canadian Arctic Archipelago and the sills between Greenland and Scotland). At no time since the inception of spreading on these northern ridges has there been a deep (thousands of meters) connection between the Arctic basins and the north Atlantic, and thus it is quite possible that endemic vent fauna of the Arctic ridges have followed an evolutionary path quite separate from any other yet studied. A return expedition planned for 2007 should answer some of these intriguing questions for the Gakkel Ridge.

The SWIR, meanwhile, occupies a crucial gap between known vent biogeographic provinces: no vent sites have yet been characterized in terms of their faunal communities between the Rodriguez Triple Junction (southern CIR, Edmond and Kairei fields) and the northern Mid-Atlantic Ridge. The CIR communities exhibit affinities with both the Atlantic and the western Pacific biogeographic provinces but are most similar overall to the western Pacific (Van Dover et al., 2001). The exploration of vent fields on the southern MAR (e.g., (German et al., 2005; Haase and M64/1-Scientific-Party, 2005)) and the SWIR thus holds promise for exciting new discoveries regarding vent organism ecology and dispersal.

Summary

Ultraslow spreading ridges have recently emerged as a unique new class of mid-ocean ridge spreading centers. Observations made on these ridges have brought tremendous advances in understanding the tectonics, magmatic architecture and hydrothermal evolution of oceanic crust at all spreading rates, and are continuing to do so. In addition, there is a growing realization that ultraslow spreading ridges provide an analog to proc-

esses that govern the breakup of continents along nonvolcanic rifted margins. The ultraslow spreading ridge community has now recognized several near-term goals on which to focus its activities, beginning with two workshops/conferences held in Europe in the fall of 2006. There will be a major effort to push ridge mapping and sampling at the slowest spreading rates into new regions, and using new technologies that mitigate both the effect of permanent ice cover, and the thick sediments that cover the Gakkel Ridge at the slowest spreading rates near the Siberian continent. In addition, many observers have recently had their eyes opened to the many ultraslow spreading ridge-relevant observations that can be made high in the Alps, where Harry Hess and Gustav Steinmann worked before them.

Acknowledgements

This article builds on the results of two conferences held on the ancient Tethyan margin of Europe to help identify and promote future priorities for the investigation of both nonvolcanic rifted margins (IODP Continental Breakup Workshop, Pontresina, Switzerland) and polar mid-ocean ridges (InterRidge Polar Ridges Meeting and Workshop, Sestri Levante, Italy). The authors thank the conference participants for their contributions, and H. Dick, W. Jokat and P. Michael for constructive reviews of this article.

References

- Aumento, F. and Loubat, H., 1970. The mid-Atlantic ridge near 45N: Serpentinized ultramafic intrusions. *Canadian Journal of Earth Sciences*, 8: 631-663.
- Bach, W., Banerjee, N.R., Dick, H.J.B. and Baker, E.T., 2002. Discovery of ancient and active hydrothermal systems along the ultra-slow spreading Southwest Indian Ridge 10°-16°E. *Geochemistry, Geophysics, Geosystems*, 3(7): 10.1029/2001GC000279.
- Baker, E.T., 1996. Geological indexes of hydrothermal venting, *Journal of Geophysical Research*, pp. 13,741-13,753.
- Baker, E.T. et al., 2004. Hydrothermal venting in magma deserts: The ultraslow-spreading Gakkel and Southwest Indian Ridges. *Geochemistry, Geophysics, Geosystems*, 5(8): Q08002, doi: 10.1029/2004GC000712.
- Baker, E.T. et al., 1994. Hydrothermal plumes along the East Pacific Rise, 8°40' to 11°50'N: Plume distribution and relationship to the apparent magmatic budget. *Earth and Planetary Science Letters*, 128: 1-17.
- Baker, E.T. and German, C.R., 2004. *Mid-Ocean Ridges: Hydrothermal Interactions Between the Lithosphere and Oceans*
On the global distribution of hydrothermal vent fields. In: C.R. German, J. Lin and L.M. Parson (Editors), *Mid-Ocean Ridges: Hydrothermal interactions between the lithosphere and oceans*. Geophysical Monograph. American Geophysical Union, Washington, DC, pp. 245-266.
- Baker, E.T. and Milburn, H.B., 1997. MAPR: A new instrument for hydrothermal plume mapping. *Ridge Events*, 8(1): 23-25.
- Bown, J.W. and White, R.S., 1994. Variation with spreading rate of oceanic crustal thickness and geochemistry. *Earth and Planetary Science Letters*, 121(3-4): 435-449.
- Cann, J. et al., 1997. Corrugated slip surfaces formed at ridge-transform intersections on the Mid-Atlantic Ridge. *Nature*, 385(6614): 329-332.
- Cannat, M. et al., 1995. Thin crust, tectonic exposures, and rugged faulting patterns at the Mid-Atlantic Ridge (22-24 Degrees North). *Geology*, 23: 49-52.
- Cannat, M., Rommevaux-Jestin, C., Sauter, D., Deplus, C. and Mendel, V., 1999. Formation of the axial relief at the very slow spreading Southwest Indian Ridge (49 degrees to 69 degrees E). *Journal of Geophysical Research - Solid Earth*, 104(B10): 22825-22843.
- Cannat, M. et al., 2006. Modes of seafloor generation at a melt-poor ultraslow-spreading ridge. *Geology*, 34(7): 605-608.
- Coakley, B. and Cochran, J., 1998. Gravity evidence of very thin crust at the Gakkel Ridge (Arctic Ocean). *Earth and Planetary Science Letters*, 162: 81-95.
- Cochran, J.R., 2003. The Gakkel Ridge; bathymetry, gravity anomalies, and crustal accretion at extremely slow spreading rates. In: G.J. Kurras, M.H. Edwards and B.J. Coakley (Editors), *Journal of Geophysical Research*. American Geophysical Union : Washington, DC, United States, United States.
- Connelly, D.P. and German, C., 2002. Geochemical anomalies over the Knipovich Ridge: Evidence for hydrothermal activity. *Eos, Transactions AGU*, 83: Ocean Sci. Meet. Suppl., Abstract OS31F-105.
- Dick, H.J.B., 1989. Abyssal peridotites, very slow spreading ridges and ocean ridge magmatism. *Geological Society Special Publications*, 42: 71-105.
- Dick, H.J.B., Fisher, R.L. and Bryan, W.B., 1984. Mineralogic variability of the uppermost mantle along mid-ocean ridges. *Earth and Planetary Science Letters*, 69(1): 88-106.

- Dick, H.J.B., Lin, J. and Schouten, H., 2003. Ultra-Slow Spreading - A New Class of Ocean Ridge. *Nature* (London), 426: 405-412.
- Edmonds, H. et al., 2003. Discovery of abundant hydrothermal venting on the ultra-slow spreading Gakkel ridge in the Arctic Ocean. *Nature* (London), 421: 252-256.
- Edwards, M.H. et al., 2001. Evidence of recent volcanic activity on the ultraslow-spreading Gakkel Ridge. *Nature*, 409(6822): 808-812.
- Engel, C.G. and Fisher, R.L., 1975. Granitic to ultramafic rock complexes of the Indian Ocean ridge system, western Indian Ocean. *Geological Society of America Bulletin*, 86(11): 1553-1578.
- Froitzheim, N. and Manatschal, G., 1996. Kinematics of Jurassic rifting, mantle exhumation, and passive-margin formation in the Austroalpine and Penninic nappes (eastern Switzerland). *Geological Society of America Bulletin*, 108(9): 1120-1133.
- German, C.R. et al., 2005. Hydrothermal activity on the southern Mid-Atlantic Ridge: tectonically- and volcanically-hosted high-temperature venting at 2-7 degrees S. *Eos, Transactions AGU*, 86(52): Fall Meet. Suppl., Abstract OS21C-04.
- Haase, K. and M64/1-Scientific-Party, 2005. Hydrothermal activity and volcanism on the southern Mid-Atlantic Ridge. *Eos, Transactions AGU*, 86(52): Fall Meet. Suppl., Abstract OS21C-04.
- Hart, S.R., 1984. A Large-Scale Isotope Anomaly in the Southern Hemisphere Mantle. *Nature*, 309(5971): 753-757.
- Hellebrand, E. and Snow, J., 2003. Deep melting underneath the highly oblique-spreading Lena Trough (Arctic Ocean). *Earth and Planetary Science Letters*, 216: 283-299.
- Hess, H.H., 1962. History of ocean basins. *Buddington Vol. (Geol. Soc. Am.)*. 599-620.
- InterRidge-Arctic-Working-Group, 1998. Mapping and Sampling Arctic Ridges: A Project Plan. In: C. Wilson (Editor), *Interridge Workshop: Mapping and Sampling the Arctic Ridges*. InterRidge, Hannover, Germany.
- Jackson, H.R., I., R. and Falconer, R.K.H., 1982. Crustal structure near the Arctic Mid-Ocean Ridge. *Journal of Geophysical Research*, 87: 1773-1783.
- Johnson, K.T.M., Dick, H.J.B. and Shimizu, N., 1990. Melting in the oceanic upper mantle; an ion microprobe study of diopsides in abyssal peridotites. *Journal of Geophysical Research, B, Solid Earth and Planets*, 95(3): 2661-2678.
- Jokat, W. et al., 2003. Geophysical evidence for reduced melt production on the super-slow Gakkel Ridge (Arctic Ocean). *Nature* (London), 423: 962-965.
- Karson, J.a.D., H., 1983. Tectonics of Ridge-Transform Intersections at the Kane Fracture Zone. *Marine Geophysical Researches*, 6(1): 51-98.
- Kelley, D.S. et al., 2001. An off-axis hydrothermal vent field near the Mid-Atlantic Ridge at 30°N. *Nature*, 412: 145-149.
- le Roex, A.P., Dick, H.J.B. and Watkins, R.T., 1992. Petrogenesis of anomalous K-enriched MORB from the Southwest Indian Ridge; 11 degrees 53'E to 14 degrees 38'E. *Contributions to Mineralogy and Petrology*, 110(2-3): 253-268.
- Manatschal, G., 2004. New models for evolution of magma-poor rifted margins based on a review of data and concepts from West Iberia and the Alps. *International Journal of earth Sciences*, 93: 432-466.
- Meyzen, C.M., Toplis, M.J., Humler, E., Ludden, J. and Mevel, C., 2003. A discontinuity in mantle composition beneath the Southwest Indian Ridge(*Nature*, 421(6974): 731-733.
- Michael, P.J. et al., 2003. Magmatic and amagmatic seafloor spreading at the slowest mid-ocean ridge: Gakkel Ridge, Arctic Ocean. *Nature* (London), 423: 956-961.

- Morgan, W.J., 1968. Rises, trenches, great faults and crustal blocks. *Journal of Geophysical Research*, 73(6): 1959-1982.
- Nicolas, A., 1995. *The Mid-Oceanic Ridges: Mountains Below Sea Level*. Springer Verlag, 200 pp.
- Okino, K. et al., 2002. Preliminary analysis of the Knipovitch Ridge segmentation, influence of focused magmatism and ridge obliquity on an ultraslow spreading system. *Earth and Planetary Science Letters*, 202: 275-288.
- Patriat, P., Sauter, D., Munsch, M. and Parson, L.M., 1997. A survey of the Southwest Indian Ridge axis between Atlantis II fracture zone and the Indian Ocean triple junction: Regional setting and large scale segmentation. *Marine Geophysical Researches*, 19(6): 457-480.
- Pedersen, R.B. et al., 2005. Two Vent Fields Discovered at the Ultraslow Spreading Arctic Ridge System. *Eos, Transactions AGU*, 86(52): Fall Meet. Suppl., Abstract OS21C-01.
- Penrose-participants, 1973. Penrose field conference on ophiolites. *Geotimes*, 17(24-25).
- Reid, I. and Jackson, H.R., 1981. Oceanic spreading rate and crustal thickness. *Marine Geophysical Researches*, 5: 165-172.
- Rona, P.A., Klinkhammer, G., Nelson, T.A., Trefry, J.H. and Elderfield, H., 1986. Black smokers, massive sulphides and vent biota at the Mid-Atlantic Ridge. *Nature*, 321: 33-37.
- Snow, J., Jokat, W., Hellebrand, E., Mühe, R. and Party, S.S., 2001a. Magmatic and hydrothermal activity in Lena Trough, Arctic Ocean. *Eos, Transactions AGU*, 82: 193-198.
- Snow, J., Jokat, W., Hellebrand, E. and Mühe, R., 2001b. Magmatic and Hydrothermal activity in Lena Trough, Arctic Ocean. *EOS Trans. AGU.*, 82: 193-198.
- Snow, J. and Petrology-Group-ARK-XX-2, 2006. Petrology of Lena Trough and Gakkel Ridge. *Reports on Polar and Marine Research*, in press.
- Steinmann, G., 1905. *Geologische Beobachtungen in den Alpen, II. Die Schardtsche Ueberfaltungstheorie und die geologische Bedeutung der Tiefseeabsätze und der ophiolithischen Massengesteine*. *Berichte Naturforschenden Gesellschaft Freiburg*, 16: 1-49.
- Tucholke, B.E., Lin, J. and Kleinrock, M., 1998. Megamullions and mullion structure defining oceanic metamorphic core complexes on the mid-Atlantic ridge. *Journal of Geophysical Research*, 103(B5): 9857-9866.
- Tyler, P.A. and Young, C.M., 2003. Dispersal at hydrothermal vents: a summary of recent progress. *Hydrobiologia*, 503: 9-19.
- Van Dover, C.L., German, C.R., Speer, K.G., Parson, L.M. and Vrijenhoek, R.C., 2002. Evolution and biogeography of deep-sea vent and seep invertebrates. *Science*, 295: 1253-1257.
- Van Dover, C.L. et al., 2001. Biogeography and ecological setting of Indian Ocean hydrothermal vents. *Science*, 294: 818-823.
- Vogt, P., Taylor, P.T., Kovacs, L.C. and Johnson, G., 1979. Detailed aeromagnetic investigation of the Arctic Basin. *Journal of Geophysical Research*, 84(B3): 1071-1089.
- Whitmarsh, R., Sawyer, D., Klaus, A. and Masson, D., 1996. *Scientific Results, Leg 149. Proceedings of the Ocean Drilling Program, Scientific Results*, 149.
- Whitmarsh, R.B., Manatschal, G. and Minshull, T.A., 2001. Evolution of magma-poor continental margins from final rifting to seafloor spreading. *Nature*, 413: 150-154.
- Whitmarsh, R.B., Pinheiro, L.M., Miles, P.R. and Sibuet, J.C., 1993. Thin crust at the Western Iberia Ocean-Continent-Transition and ophiolites. *Tectonics*, 12(5): 1230-1239.